

## KDQOL-36 Average Scores\*

<b>Males/Non-Diabetic</b>	<b>Age</b>	<b>Age</b>	<b>Age</b>	<b>Age</b>
<b>Variable</b>	<b>&lt;45</b>	<b>45-64</b>	<b>65-74</b>	<b>75+</b>
PCS	41	39.4	35.4	33.8
MCS	46.2	48.9	45.5	47.7
Burden of Illness	43.6	47.6	41.6	46.1
Physical Symptoms	75.1	76.9	72.6	76
Effect of Kidney Disease	62.6	67.3	64.8	68.1

<b>Males/ Diabetic</b>	<b>Age</b>	<b>Age</b>	<b>Age</b>	<b>Age</b>
<b>Variable</b>	<b>&lt;45</b>	<b>45-64</b>	<b>65-74</b>	<b>75+</b>
PCS	37.1	34.1	34.3	31.9
MCS	47.4	45.3	50.5	42.8
Burden of Illness	43.5	38	45.2	36.8
Physical Symptoms	73.2	72.6	77.1	73.2
Effect of Kidney Disease	68.8	60.3	69.0	68.8

\* Based on KDQOL-36 data from 1,282 US Prevalent in-center hemodialysis patients supplied by Arbor Research Collaborative for Health and categorized by age, gender, diabetic/non-diabetic.

## KDQOL-36 Average Scores\*

<b>Females/ Non-Diabetic</b>	<b>Age</b>	<b>Age</b>	<b>Age</b>	<b>Age</b>
<b>Variable</b>	<b>&lt;45</b>	<b>45-64</b>	<b>65-74</b>	<b>75+</b>
PCS	39.7	37	30.8	32.2
MCS	45.8	47.6	48.5	45
Burden of Illness	44.7	48.7	44.0	42.7
Physical Symptoms	72.2	70	74.8	73.6
Effect of Kidney Disease	61.1	61.3	69.4	66.6

<b>Females/ Diabetic</b>	<b>Age</b>	<b>Age</b>	<b>Age</b>	<b>Age</b>
<b>Variable</b>	<b>&lt;45</b>	<b>45-64</b>	<b>65-74</b>	<b>75+</b>
PCS	35.7	31.9	31.3	29.3
MCS	46.8	45.4	47.4	49.2
Burden of Illness	31.1	42.1	44.5	45.0
Physical Symptoms	73.1	69.9	71	75.5
Effect of Kidney Disease	54.8	63.8	65.4	70.8

\* Based on KDQOL-36 data from 1,282 US Prevalent in-center hemodialysis patients supplied by Arbor Research Collaborative for Health and categorized by age, gender, diabetic/non-diabetic.